

PubMed

Format: Abstract

Full text links



J Arthroplasty. 2017 Sep;32(9):2788-2791. doi: 10.1016/j.arth.2017.03.052. Epub 2017 Mar 31.

Cryotherapy With Dynamic Intermittent Compression Improves Recovery From Revision Total Knee Arthroplasty.

Murgier J¹, Cailliez J¹, Wargny M², Chiron P¹, Cavaignac E¹, Laffosse JM¹.

Author information

Abstract

BACKGROUND: The goal of this study was to assess the efficacy of cryotherapy with dynamic intermittent compression (CDIC) in relieving postoperative pain, decreasing blood loss, and improving functional scores after revision total knee arthroplasty (rTKA).

METHODS: We conducted a prospective case-control study (level of evidence: I) to evaluate the efficacy of CDIC on postoperative bleeding, pain, and functional outcomes after rTKA. Forty-three cases were included at a single institution and divided in 2 groups: a control group without CDIC (n = 19) and an experimental group with CDIC (n = 24). Bleeding was evaluated by calculating total blood loss, pain at rest was evaluated with a visual analog scale on postoperative day 3, and function was assessed using the Oxford score at 6 months postoperatively. The comparative analysis was performed using the Fisher exact test.

RESULTS: The CDIC group had significantly lower total blood loss (260 vs 465 mL; $P < .05$), significantly less pain on day 3 (1 vs 3; $P < .05$), and a significantly higher functional score (42 vs 40; $P < .05$) than the control group.

CONCLUSION: This is the first report dealing with the use of CDIC after rTKA. According to our results, it improves the recovery of patients who underwent rTKA; thus, it should be integrated into our daily practice.

Copyright © 2017 Elsevier Inc. All rights reserved.

KEYWORDS: arthroplasty; blood loss; compressive cryotherapy; knee; pain

PMID: 28465126 DOI: [10.1016/j.arth.2017.03.052](https://doi.org/10.1016/j.arth.2017.03.052)

[Indexed for MEDLINE]


MeSH terms

LinkOut - more resources

Accepted Manuscript

Cryotherapy with Dynamic Intermittent Compression Improves Recovery from Revision Total Knee Arthroplasty

J. Murgier, J. Cailliez, M. Wargny, P. Chiron, E. Cavaignac (, J.M. Laffosse



PII: S0883-5403(17)30289-9

DOI: [10.1016/j.arth.2017.03.052](https://doi.org/10.1016/j.arth.2017.03.052)

Reference: YARTH 55778

To appear in: *The Journal of Arthroplasty*

Received Date: 17 February 2017

Revised Date: 9 March 2017

Accepted Date: 22 March 2017

Please cite this article as: Murgier J, Cailliez J, Wargny M, Chiron P, Cavaignac (E, Laffosse J, Cryotherapy with Dynamic Intermittent Compression Improves Recovery from Revision Total Knee Arthroplasty, *The Journal of Arthroplasty* (2017), doi: [10.1016/j.arth.2017.03.052](https://doi.org/10.1016/j.arth.2017.03.052).

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Cryotherapy with Dynamic Intermittent Compression Improves Recovery from Revision Total Knee Arthroplasty

*Effectiveness of cryotherapy with dynamic intermittent compression in
rTKA*

J. Murgier* (1), J. Cailliez (1), M. Wargny (2), P. Chiron, E. Cavaignac
(1), JM. Laffosse (1)

(1) Département de chirurgie Orthopédique et Traumatologique,
CHU Toulouse, Toulouse, France

(2) Epidemiology Department, CHU Toulouse, Toulouse, France

***Corresponding author:**

Dr Murgier Jérôme

Département d'Orthopédie Traumatologie

CHU Toulouse – Hôpital Pierre Paul Riquet

Place du Docteur Baylac, TSA 40031

31059 Toulouse Cedex 9, France

Email: murgier.jerome@hotmail.fr

Tel: 0561775582

Fax:0561775432

Conflict of interest: No benefits in any form have been received or will be received
from a commercial party related directly or indirectly to the subject of this article.

Cryotherapy with Dynamic Intermittent Compression Improves Recovery from Revision Total Knee Arthroplasty

Effectiveness of cryotherapy with dynamic intermittent compression in rTKA

ABSTRACT

Purpose : The goal of this study was to Assess the efficacy of cryotherapy with dynamic intermittent compression (CDIC) in Relieving post operative pain, decreasing blood loss and improving functional scores after revision total knee arthroplasty (rTKA).

Methods: we conducted a prospective case-control study. (Level of evidence: I) to evaluate the efficacy of CDIC on postoperative bleeding, pain and functional outcomes after rTKA. 43 cases were included at a single institution and divided in two groups: a control group without CDIC (n = 19) and an experimental group with CDIC (n = 24). Bleeding was evaluated by calculating total blood loss; pain at rest was evaluated with a visual analogue scale (VAS) on postoperative day 3; function was assessed using the Oxford score at 6 months postoperative. The comparative analysis was performed using Fisher's exact test

Results: The CDIC group had significantly lower total blood loss (260 ml vs 465 ml, $P < 0.05$), significantly less pain on day 3 (1 vs 3, $P < 0.05$) and a significantly higher functional score (42 vs 40, $P < 0.05$) than the control group.

Conclusion: This is the first report dealing with the use of CDIC after rTKA. According to our results , it improves the recovery of patients who underwent revision TKA, thus it should be integrated into our daily practice.

32 INTRODUCTION

33

34 Revision total knee arthroplasty (rTKA) procedures cause blood loss. Postoperative anaemia is
35 associated with a higher risk of infection [7], of patient dissatisfaction [9] and postoperative blood transfusion
36 with its inherent risks [23].

37 Revision TKA is a major orthopaedic procedure that causes soft tissue damages which contribute to
38 localised pain, which in turn reduces range of motion and causes persistent quadriceps atrophy [25]. Significant
39 blood loss after this procedure (up to 1.5 L) can lead to systemic complications [23]. Despite progress in
40 multimodal analgesia and anaesthetic methods, knee arthroplasty is a painful surgery [25]. Non-pharmacological
41 treatments can also play a role, most notably cryotherapy which decreases the local metabolism, thereby
42 reducing blood loss and pain [13]. This technique has minimal disadvantages relative to its potential benefits
43 [11].

44 New devices that combine cryotherapy with dynamic intermittent compression (CDIC) have recently
45 been introduced. These devices provide a dry cold and maintain a consistent temperature for an extended period
46 of time [24, 25]. While the benefits of these systems were demonstrated in primary TKA [24] and anterior
47 cruciate ligament reconstruction [17], we did not find any published studies evaluating the effect of CDIC in
48 patients undergoing revision TKA.

49 Our hypothesis was that use of CDIC would reduce total blood loss after revision TKA. The main
50 objective of this study was to assess its efficacy in terms of postoperative blood loss. Patients who underwent
51 rTKA were split into two matched groups for comparisons: one group with CDIC and the other one without. The
52 other objectives were to compare the blood transfusion rate, the pain, the functional scores and the complication
53 rate in both groups.

54

55 PATIENTS AND METHODS

56
57 This was a single-institution, prospective case-control study (Level of evidence: III). It was approved by
58 our hospital's research ethics committee (Number 01-0115).

59

60 ***Patients***

61 All patients who underwent single-stage rTKA from January 2013 to January 2015 were included.

62 Patients were excluded when a two-stage revision or a partial revision was performed. They were also
63 excluded if they had a contraindication to CDIC, such as history of deep vein thrombosis, a coagulation disorder
64 or skin damage at the device application site.

65 Forty-three patients were included (27 males, 16 females). The revisions procedures were carried out
66 with a rotating hinge knee prosthesis (RHK Nexgen®, Zimmer, Warsaw, USA) in all cases. A tibial tubercle
67 osteotomy was needed in 9 cases and the patella was resurfaced in 14 cases during the revision procedure.

68 The anaesthesia and postoperative analgesia protocols used were standardised and similar in the two
69 groups. Anticoagulant therapy was initiated 6 hours after the end of surgery in all patients.

70 The procedure was performed with a tourniquet in all cases. It was released before closing the wound to
71 realize complete haemostasis. The mean procedure duration was 120 minutes (90–140).

72

73 ***Methods***

74 The population was divided into two groups: a control group without CDIC and an experimental one
75 with CDIC. The demographics data in these two groups were comparable (Table 1). The patients in the control
76 group were included between January 2013 and April 2014. The patients in the CDIC group were included
77 between May 2014 and January 2015.

78 The CDIC device used was the Game Ready® system (CoolSystems Inc., Concord, CA, USA). It
79 comes with an anatomical wrap that is applied to the knee. This wrap circulates pre-cooled compressed air and
80 water. The temperature-controlled unit generates a dry cold; this is more comfortable for the patient than wet
81 cold, thereby limiting the risk of maceration, bandage deterioration and skin lesions. The wrap is covered with a
82 removable, washable cover for the patient's health and comfort. The wrap is connected to a portable control unit.
83 The compression is applied intermittently depending on the protocol selected. The surgery support staff was
84 given specific training on how to use the CDIC.

85 The following protocol was used:

- 86 - Application: after bandaging, in the operating room and before transfer to recovery room
- 87 - Intensity: programme 3 (30 minute on/off cycles)
- 88 - Temperature: 8°C
- 89 - Application duration: two 8-hour cycles over a 24-hour period
- 90 - Treatment duration: 72 hours postoperative

91 The control group was treated with regular cold application (4 hours per day) using a cold pack.

92 The following parameters were measured in both groups: total blood loss, haemoglobin and haematocrit
 93 levels on D-1, D+1 and D+5, transfusion volume and rate (red cell concentrate (RCC) units), pain on
 94 postoperative day 3, functional outcomes based on the Oxford score at 6 months postoperative and the number of
 95 complications recorded at 6 months postoperative.

96 Total blood loss was calculated using the preoperative (D-1) and postoperative (D+5) laboratory test
 97 results according to the Mercuriari formula [8]:

98 Total blood loss = $VST \times (Hct_{pre} - Hct_{post\ D5}) + \text{volume of retransfused RCC}^*$

99 where the patient's total blood volume = $k_1 \times \text{height (m)}^3 + k_2 \times \text{mass (kg)} + k_3$

100 for men: $k_1 = 0.3669$, $k_2 = 0.03219$, and $k_3 = 0.6041$;

101 and for women: $k_1 = 0.3561$, $k_2 = 0.03308$, and $k_3 = 0.1833$

102 Hct_{pre} = initial preoperative Hct

103 $Hct_{post\ D5}$ = Hct on the morning of the 5th postoperative day

104 When transfusion was done (allogenic or autologous), the total blood loss was equal to the blood loss calculated
 105 from the change in haematocrit plus the volume transfused [14].

106 The indication for RCC transfusion in our surgical unit is standardised to Hb < 8 g/dL and/or patient
 107 with symptomatic anaemia.

108 Postoperative pain at rest was measured by the surgery unit's nurse using a visual analogue scale (VAS)
 109 on the 3rd day postoperative and by looking at the cumulative morphine use on the 5th postoperative day,
 110 expressed in morphine-equivalent dose (in mg).

111 The Oxford score [4] was collected pre- and postoperatively using the validated French version of the
 112 questionnaire [5]. The questionnaire was filled out during a follow-up visit 6 months after the procedure.

113

114 **Statistical analysis**

115 Cohort characteristics are presented as numbers, means, SDs, and ranges. The normal distribution of the
 116 data was assessed using the Kolmogorov–Smirnov test. For variables that were not normally distributed, data
 117 were analysed using the Mann-Whitney test for independent samples and the Wilcoxon signed rank test for
 118 dependent samples. Comparison of observed proportions was performed using Fisher's exact test. Statistical
 119 analysis was carried out using SPSS 18 Statistical Software (SPSS Inc, Chicago, IL, USA) and significance was
 120 set at P less than 0.05.

121

122 **RESULTS (Table2)**

123

124 **Bleeding**

125 The total blood loss was lower in the CDIC group than in the control group (260 ml vs 465 ml, $P <$
 126 0.05). The haemoglobin and haematocrit levels were similar between groups. The transfusion rate was lower in
 127 the CDIC group (8% vs 42%, $P < 0.05$) and the mean lowest Haemoglobin level was lower in the control group
 128 with 8.5 gm/dL (+/- 1,2) vs 9,6 (+/- 1,6) ; $p < 0,005$. In the CDIC group, the number of RCC units given per
 129 patient was lower as well. No differences were found in any of the other measured blood-related parameters.

130

131 Pain

132 Pain at rest on day 3 was lower in the CDIC group than in the control group (1 vs 3, $P < 0.05$). The
133 cumulative morphine intake at day 5 was not significantly different between groups.

134

135 Functional scores at 6 months

136 The Oxford score at 6 months postoperative was higher in the CDIC group than in the control group (42
137 vs 40).

138

139 Complications

140 There were four complications in the CDIC group and three in the control group. There were two cases
141 of infection recurrence and one case of deep vein thrombosis in each group, and one case of extensor mechanism
142 disruption in the CDIC group.

143

144

145 DISCUSSION

146

147 Our hypothesis was confirmed. The patients in the CDIC group had lower total blood loss than patients
148 in the control group. Moreover, the transfusion rate and the pain were lower in the CDIC group. The functional
149 outcome was similar between the two groups at 6 months postoperative.

150 This is the first study to evaluate the use of CDIC after revision TKA. The blood loss was evaluated
151 using a method previously validated for revision TKA patients [22]. This method provides a complete view of
152 the total blood loss, as it also takes into account hidden blood loss following TKA [23]. We did not use the blood
153 volume present in the suction drains, as in other studies [16]. This blood loss calculation method is not reliable;
154 it overestimates blood losses and can lead to more blood transfusions [19]. The volume of blood in surgical
155 drains has never been validated as being an objective measure of blood loss [22]. There is no correlation between
156 the volume of blood in the drains and the need for transfusion [15].

157 CDIC has been used in the sports medicine setting to improve recovery and to treat ligament and bone
158 injuries [15, 17]. It has been shown that CDIC improves postoperative recovery by stimulating the tissue repair
159 process [3]. Generally, these systems are provided to healthy athletes undergoing a minor procedure compared to
160 revision TKA. We believe that any help is beneficial to a fragile population such as the one undergoing revision
161 TKA.

162 The contribution of CDIC to postoperative recovery from TKA has already been demonstrated. Su et al
163 [24] evaluated CDIC in patients undergoing primary TKA and compared it to a control group. In that study, the
164 patients in the CDIC group used the system for 5 days after the procedure. They found a lower narcotic intake
165 and slight improvement in the functional outcome in the patients using CDIC. We also found a tendency of
166 reduced narcotic use (-20 mg morphine-equivalents in the CDIC group). This reduced narcotic intake reduces

167 the side effects inherent to these agents. Patients feel less medicated and have a better postoperative course.
168 Advanced cryotherapy was compared to icing only in the postoperative course of TKA in a randomised
169 controlled trial [25]. The authors concluded that there were no advantages in using advanced cryotherapy in
170 daily practice, particularly because of the additional cost associated with these systems. However, more than one
171 kind of cryotherapy system was used in that study, leading to variability in the results. Moreover, the blood loss
172 was measured only through haemoglobin variations, which does not take into account hidden blood loss [23].
173 Additionally, no blood transfusions were performed, as the patients were undergoing primary TKA. The need for
174 transfusion is higher during revision TKA [2].

175 Other therapeutic means have been proposed to reduce bleeding during primary and revision TKA
176 procedures. Tranexamic acid has been shown to be effective in hip and knee arthroplasty [1]. Use of thrombin-
177 based topical haemostatics does not have clear-cut benefits. One group has described its benefits in revision
178 TKA [22]; however, anaemia, atrial fibrillation, infection have been associated with this type of product [12, 16,
179 18, 21]. These side effects do not come into play when using CDIC. Thienpont et al [25] bring up the risk of
180 frostbite in the area where CDIC is applied. This is an extremely serious complication that would require an
181 additional major soft tissue procedure [6]. We have not encountered this complication, and have not found any
182 documented cases of frostbite with CDIC.

183
184 The current study has certain limitations. Firstly, this was a multi-surgeon study which increases the
185 variability of the results. However, this also means that the study can be more easily generalised to current
186 practice. Secondly, this study was performed within a highly specialised TKA surgery unit. Because of the use of
187 advanced anaesthesia procedures, analgesic infiltration and preventative multimodal pain management [11], it is
188 possible that a type II error occurred in our interpretation of the results. However, the anaesthesia and analgesia
189 techniques did not differ between the control and CDIC groups. Moreover, the transfusion rate – likely the most
190 relevant criteria from a clinical point of view – is subjected to confounding factors because of the patients' co-
191 morbidities. Although we use a standardised approach, this bias is still present.

192 Conclusion

193
194
195 The number of revision TKA procedures performed each year will continue to increase [10, 20, 22].
196 Since CDIC improves the recovery of patients undergoing revision TKA, it should be integrated into our daily
197 practice. Prospective randomised trial is necessary to validate the results of our study.

198
199
200
201
202
203
204
205

206

207

208

REFERENCES

- 209 1. Alshryda S, Sukeik M, Sarda P, et al (2014) A systematic review and meta-analysis of
210 the topical administration of tranexamic acid in total hip and knee replacement. *Bone*
211 *Joint J* 96-B:1005–1015.
- 212 2. Berman AT, Geissele AE, Bosacco SJ (1988) Blood loss with total knee arthroplasty.
213 *Clin Orthop Relat Res* 137–138.
- 214 3. Dahl J, Li J, Bring DK-I, et al (2007) Intermittent pneumatic compression enhances
215 neurovascular ingrowth and tissue proliferation during connective tissue healing: a
216 study in the rat. *J Orthop Res* 25:1185–1192.
- 217 4. Dawson J, Fitzpatrick R, Carr A, Murray D (1996) Questionnaire on the perceptions of
218 patients about total hip replacement. *J Bone Joint Surg Br* 78:185–190.
- 219 5. Delaunay C, Epinette J-A, Dawson J, et al (2009) Cross-cultural adaptations of the
220 Oxford-12 HIP score to the French speaking population. *Orthop Traumatol Surg Res*
221 95:89–99.
- 222 6. Dundon JM, Rymer MC, Johnson RM (2013) Total patellar skin loss from cryotherapy
223 after total knee arthroplasty. *J Arthroplasty* 28:376.e5–7.
- 224 7. Freedman J, Luke K, Monga N, et al (2005) A provincial program of blood
225 conservation: The Ontario Transfusion Coordinators (ONTraC). *Transfus Apher Sci*
226 33:343–349.
- 227 8. Gibon E, Courpied J-P, Hamadouche M (2013) Total joint replacement and blood loss:
228 what is the best equation? *Int Orthop* 37:735–739.
- 229 9. Ishida K, Tsumura N, Kitagawa A, et al (2011) Intra-articular injection of tranexamic
230 acid reduces not only blood loss but also knee joint swelling after total knee
231 arthroplasty. *Int Orthop* 35:1639–1645.
- 232 10. Jeserschek R, Clar H, Aigner C, et al (2003) Reduction of blood loss using high-dose
233 aprotinin in major orthopaedic surgery: a prospective, double-blind, randomised and
234 placebo-controlled study. *J Bone Joint Surg Br* 85:174–177.
- 235 11. Kehlet H, Thienpont E (2013) Fast-track knee arthroplasty -- status and future
236 challenges. *Knee* 20 Suppl 1:S29–33.
- 237 12. Kim HJ, Fraser MR, Kahn B, et al (2012) The efficacy of a thrombin-based hemostatic
238 agent in unilateral total knee arthroplasty: a randomized controlled trial. *J Bone Joint*
239 *Surg Am* 94:1160–1165.
- 240 13. Kullenberg B, Ylipää S, Söderlund K, Resch S (2006) Postoperative cryotherapy after
241 total knee arthroplasty: a prospective study of 86 patients. *J Arthroplasty* 21:1175–
242 1179.

- 243 14 Liu X, Zhang X, Chen Y, et al (2011) Hidden blood loss after total hip arthroplasty. *J*
244 *Arthroplasty* 26:1100–5.e1. doi: 10.1016/j.arth.2010.11.013
- 245 15 Mesa-Ramos F, Mesa-Ramos M, Maquieira-Canosa C, Carpintero P (2008) Predictors
246 for blood transfusion following total knee arthroplasty: a prospective randomised
247 study. *Acta Orthop Belg* 74:83–89.
- 248 16. Mozet C, Prettin C, Dietze M, et al (2012) Use of Floseal and effects on wound healing
249 and pain in adults undergoing tonsillectomy: randomised comparison versus
250 electrocautery. *Eur Arch Otorhinolaryngol* 269:2247–2254.
- 251 17 Murgier J, Cassard X (2014) Cryotherapy with dynamic intermittent compression for
252 analgesia after anterior cruciate ligament reconstruction. Preliminary study. *Orthop*
253 *Traumatol Surg Res* 100:309–312.
- 254 18. Nasso G, Piancone F, Bonifazi R, et al (2009) Prospective, randomized clinical trial of
255 the FloSeal matrix sealant in cardiac surgery. *Ann Thorac Surg* 88:1520–1526.
- 256 19. Parker MJ, Livingstone V, Clifton R, McKee A (2007) Closed suction surgical wound
257 drainage after orthopaedic surgery. *Cochrane Database Syst Rev* CD001825.
- 258 20. Rasouli MR, Harandi AA, Adeli B, et al (2012) Revision total knee arthroplasty:
259 infection should be ruled out in all cases. *J Arthroplasty* 27:1239–43.e1–2.
- 260 21. Renkens KL, Payner TD, Leipzig TJ, et al (2001) A multicenter, prospective,
261 randomized trial evaluating a new hemostatic agent for spinal surgery. *Spine* 26:1645–
262 1650.
- 263 22. Romanò CL, Monti L, Logoluso N, et al (2014) Does a thrombin-based topical
264 haemostatic agent reduce blood loss and transfusion requirements after total knee
265 revision surgery? A randomized, controlled trial. *Knee Surg Sports Traumatol*
266 *Arthrosc.*
- 267 23. Sehat KR, Evans RL, Newman JH (2004) Hidden blood loss following hip and knee
268 arthroplasty. Correct management of blood loss should take hidden loss into account. *J*
269 *Bone Joint Surg Br* 86:561–565.
- 270 24. Su EP, Perna M, Boettner F, et al (2012) A prospective, multi-center, randomised trial
271 to evaluate the efficacy of a cryopneumatic device on total knee arthroplasty recovery.
272 *J Bone Joint Surg Br* 94:153–156.
- 273 25. Thienpont E (2014) Does advanced cryotherapy reduce pain and narcotic consumption
274 after knee arthroplasty? *Clin Orthop Relat Res* 472:3417–3423.

275

276

277

Table 1 Baseline characteristics and surgery related data for the cohort
 TT: tibial tubercle, BMI: Body mass index, ASA: American Society of Anesthesiologists

		Control group (n=19)	CDIC group (n=24)	P
Baseline characteristics	Mean (SD) Age, years	66.5 (9.7)	70 (13.9)	0.86
	Sex ratio : F/M	8/11	8/16	0.78
	Mean (SD) BMI, kg/m ²	29.7 (4.6)	29.7 (4.7)	0.89
	Pré op anticoagulants, n (%)	6 (32)	3 (13)	0.15
	ASA score, n (%)	1: 1 (5) 2: 8 (42) 3: 10 (53)	1: 4 (17) 2: 14 (58) 3: 6 (25)	0.17
Surgery-related data	Mean (SD) Surgery time, min	120 (40.2)	118 (43.3)	0.84
	Mean (SD) Tourniquet time	98 (27.4)	100 (29.3)	0.98
	TT osteotomy, n (%)	5 (26)	4 (17)	0.48
	Patella resurfacing, n (%)	8(42)	6(25)	0.39

Table 2 Summary of variables measured in both groups.

RCC: red cell concentrate, Hb: haemoglobin, Hct: haematocrit, VAS: visual analogue scale

		Control group (n=19)	CDIC group (n=24)	P
Blood loss	Mean (SD) Hb at D-1 (g/dL)	12.5 (2.1)	13 (1.8)	0.76
	Mean (SD) Hb at D+5 (g/dL)	10.4 (1.2)	10.7 (1.3)	0.31
	Mean (SD) Hct at D-1 (%)	37 (5.9)	39.6 (4.9)	0.9
	Mean (SD) Hct at D+5 (%)	31 (3.6)	32.7 (3.6)	0.19
	Mean (SD) Total blood loss (ml)	465 (275)	260 (106)	0.024
	Mean lowest Hb level	8.5	9.6	0.03
	Transfusion rate	42%	8%	0.013
Transfusion	Number of RCC units	0: 58% (n=11) 1: 5% (n=1) 2: 26% (n=5) 3: 11% (n=2)	0: 92% (n=22) 1: 0% (n = 0) 2: 4% (n=1) 3: 4% (n=1)	0.023
Pain	Mean (SD) VAS Day +3	3 (1)	1(1)	0.01
	Mean (SD) Narcotic consumption at Day +5 (mg)	100 (37)	80 (37)	1
Functional score	Mean (SD) Oxford	40 (2.8)	42 (2.4)	NS